

APPENDIX D: DISASTER/EMERGENCY PREPAREDNESS

DISASTER/EMERGENCY NUMBERS	CONTACT/TOWN	TELEPHONE NUMBER
AMBULANCE		
APPLIANCE REPAIR		
BUILDING INSPECTOR		
CHILD ABUSE HOT LINE		1-800-482-5964
CHILD CARE LICENSING UNIT	Little Rock	1-800-445-3316 or 501-682-8590
CLEANING and MAINTENANCE		
ELECTRIC COMPANY		
ELECTRICIAN		
EMERGENCY CHILD LOCATOR		1-866-908-9572
FACILITY DIRECTOR		
FIRE DEPARTMENT		
FIRE DEPARTMENT (Non-Emergency)		
GAS COMPANY		
GLASS COMPANY		
HEALTH DEPARTMENT (Local)		
HEATING and AIR CONDITIONING		
INSURANCE AGENT AND POLICY NUMBER		
LICENSING SPECIALIST		
LOCKS		
NATIONAL EMERGENCY FAMILY REGISTRY AND LOCATOR PLUMBER		1-800-588-9822
POISON CONTROL		1-800-376-4766
POLICE		
POLICE (Local Non-Emergency)		
RED CROSS (Local)		
SHERIFF		
TRASH REMOVAL		
WATER DEPARTMENT		

IMMUNIZATION REQUIREMENTS

Table I:

Immunization Requirements for Child Care and Early Childhood Education Facilities

Instructions for utilizing Table I: Table I is not a recommendation of vaccines to get, but of doses required to already have at that age. To determine what vaccines are required for a child to attend a licensed child care facility, refer to Column one (1) on the left to see what age range is correct for the child. Then all the vaccines on the same row as the child's age are required for attendance in a licensed child care facility. Vaccines are required based on the current age of the child. Column one (1) is not an age range for when a child can be vaccinated.

<u>Column 1</u>	<u>Column 2</u>	<u>Column 3</u>	<u>Column 4</u>	<u>Column 5</u>	<u>Column 6</u>	<u>Column 7</u>	<u>Column 8</u>	<u>Column 9</u>
Current AGE of child	DTaP DTP/DT	POLIO	Hib **	HEPATITIS B	MMR ****	VARICELLA ****	PNEUMOCOCCAL **	HEPATITIS A
1-2 Months	None	None	None	None (1-2 doses possible)	None	None	None	
3-4 Months	1 dose	1 dose	1 dose	1 dose (1-2 doses possible)	None	None	1 dose	
5-6 Months	2 doses OR 1 dose within last 8 weeks	2 doses OR 1 dose within last 8 weeks	2 doses OR 1 dose within last 8 weeks	2 doses OR 1 dose within last 8 weeks	None	None	2 doses OR 1 dose within last 8 weeks	
7-12 Months	3 doses OR 1 dose within last 8 weeks	2 doses OR 1 dose within last 8 weeks (3 doses possible)	2-3 doses OR 1 dose within last 8 weeks	2 doses OR 1 dose within last 8 weeks (3 doses possible)	None	None	2-3 doses OR 1 dose within last 8 weeks	
13-15 Months	3 doses OR 1 dose within last 8 weeks	2 doses OR 1 dose within last 8 weeks (3 doses possible)	2-3 doses OR 1 dose within last 8 weeks (4 doses possible)	2 doses OR 1 dose within last 8 weeks (3 doses possible)	None (1 dose possible)	None (1 dose possible. A medical professional history of disease may be accepted in lieu of receiving vaccine.)	2-3 doses OR 1 dose within last 8 weeks (4 doses possible)	
16-18 Months	3 doses or 1 dose within last 8 weeks	2 doses or 1 dose within last 8 weeks (3 doses possible)	3-4 doses with last dose on/after 1 st birthday OR 2 doses if first dose is administered at age 12 - 14 months and doses are at least 8 weeks apart OR 1 dose on/after 15 months of age if no prior doses	2 doses OR 1 dose within the last 8 weeks (3 doses possible)	1 dose	1 dose A medical professional history of disease may be accepted in lieu of receiving vaccine.	3-4 doses with last dose must be on/after 1 st birthday OR 2 doses on/after 1 st birthday	

<u>Column 1</u>	<u>Column 2</u>	<u>Column 3</u>	<u>Column 4</u>	<u>Column 5</u>	<u>Column 6</u>	<u>Column 7</u>	<u>Column 8</u>	<u>Column 9</u>
Current AGE of child	DTaP DTP/DT	POLIO	Hib **	HEPATITIS B	MMR ****	VARICELLA ****	PNEUMOCOCCAL **	HEPATITIS A
19-48 months	4 doses OR 3rd dose within last 6 months OR 1 dose within last 8 weeks	3 doses OR 1 dose within last 8 weeks	3-4 doses with last dose on/after 1 st birthday OR 2 doses if first dose is administered at age 12 - 14 months and doses are at least 8 weeks apart OR 1 dose on/after 15 months of age if no prior doses	3 doses *** OR 1 dose within last 8 weeks	1 dose	1 dose A medical professional history of disease may be accepted in lieu of receiving vaccine.	3-4 doses with last dose must be on/after 1st birthday OR 1 dose on/after 24 months of age if no prior doses OR 2 doses on/after 1 st birthday	For 19-24 months: 1 dose on or after first birthday (2 doses possible) For 25-48 months: 2 doses with one dose on or after 1st birthday and at least 6 months from first dose
≥49 months	5 doses * OR 4 th dose within last 6 months OR 1 dose within last 8 weeks OR 4 doses with last dose on/after 4 th birthday	4 doses with a minimum interval of 6 months between the 3 rd and 4 th dose OR 1 dose within last 8 weeks	3-4 doses with last dose on/after 1 st birthday OR 2 doses if first dose is administered at age 12 - 14 months and doses are at least 8 weeks apart OR 1 dose on/after 15 months of age if no prior doses Not required on/after 5th birthday	3 doses *** OR 1 dose within the last 8 weeks	1 dose	1 dose A medical professional history of disease may be accepted in lieu of receiving vaccine.	3-4 doses with last dose on/after 1 st birthday OR 1 dose on/after 24 months of age if no prior doses OR 2 doses on/after 1 st birthday Not required on/after 5th birthday	2 doses with one dose on or after 1st birthday and at least 6 months from first dose

*5th DTaP/DTP/DT (Pre-school dose) must be given on or after the child's 4th birthday. Interval between 4th DTaP/DTP/DT and 5th DTaP/DTP/DT should be at least six (6) months. If a child is currently \geq forty-nine (49) months of age and does not meet the above criteria or is in process within fifteen (15) days, they are not up-to-date and should be scheduled for immunization.

** For Hib and Pneumococcal, children receiving the first dose of vaccine at age seven (7) months or older require fewer doses to complete the series.

*** 3rd dose of hepatitis B should be given at least eight (8) weeks after the 2nd dose, at least sixteen (16) weeks after the 1st dose, and it should not be administered before the child is twenty-four (24) weeks of age.

**** Vaccine doses administered up to four (4) days before the minimum interval or minimum age can be counted as valid for doses already administered. Exception: The minimum interval between doses of live vaccines (such as MMR and Varicella) must be twenty-eight (28) days.

*****A medical professional is a medical doctor (MD), advanced practice nurse (APN), doctor of osteopathy (DO), or physician assistant (PA). No self or parental history of disease will be accepted.

TABLE II: KINDERGARTEN THROUGH GRADE TWELVE (12) IMMUNIZATION REQUIREMENTS*

Vaccine ► ----- Grade ▼	Diphtheria, Tetanus, Pertussis (DTP/DT/Td/DTaP/ Tdap)	Polio (OPV – Oral or IPV – Inactivated)	MMR**** * (Measles, Mumps, and Rubella)	Hep B	Meningococcal (MCV4)	Varicella	Hepatitis A
Kindergarten	4 doses (with 1 dose on or after 4 th birthday)	3 doses (with 1 dose on or after 4 th birthday and a minimum interval of 6 months between the 2 nd and 3 rd dose) OR 4 doses with 1 dose on or after 4 th birthday and a minimum interval of 6 months between the 3 rd and 4 th dose	2 doses (with dose 1 on or after 1 st birthday and dose 2 at least 28 days after dose 1)	3 doses	None	2 doses (with dose 1 on or after 1 st birthday and dose 2 at least 28 days after dose 1) *****A medical professional history of disease may be accepted in lieu of receiving vaccine.	1 dose on or after 1 st birthday

Vaccine ► Grade ▼	Diphtheria, Tetanus, Pertussis (DTP/DT/Td/DTaP/Tdap)	Polio (OPV – Oral or IPV – Inactivated)	MMR**** * (Measles, Mumps, and Rubella)	Hep B	Meningococcal (MCV4)	Varicella	Hepatitis A
Grades 1 – 12	<p>4 doses (with 1 dose on or after 4th birthday)</p> <p>AND</p> <p>1 dose of Tdap for ages 11 years (as of September 1st each year) and older</p> <p>OR</p> <p>3 doses***** for persons 7 years of age or older who are not fully vaccinated (including persons who cannot document prior vaccination)</p>	<p>3 doses (with 1 dose on or after 4th birthday with a minimum interval of 6 months between the 2nd and 3rd dose)</p> <p>OR</p> <p>4 doses with 1 dose on or after 4th birthday and a minimum interval of 6 months between the 3rd and 4th dose</p>	<p>2 doses (with dose 1 on or after 1st birthday and dose 2 at least 28 days after dose 1)</p>	<p>2** or 3*** doses (11-15 year olds could be on a 2-dose schedule)</p>	<p>Second dose at age 16 years (as of September 1st each year) with a minimum interval of 8 weeks since 1st dose</p> <p>OR</p> <p>1 dose if not vaccinated prior to age 16 years (If first dose is administered at age 16 years or older, no second dose required.)</p>	<p>2 doses (with dose 1 on or after 1st birthday and dose 2 at least 28 days after dose 1)</p> <p>OR</p> <p>*****A medical professional history of disease may be accepted in lieu of receiving vaccine.</p>	<p>Grade 1 only: 1 dose on or after 1st birthday</p>

*Doses of vaccine required for school entry may be less than the number of doses required for age-appropriate immunization.

**An alternative two (2) dose hepatitis B schedule for eleven (11) – fifteen (15) year-old children may be substituted for the three (3) dose schedule. Only an FDA-approved alternative regimen vaccine for the two (2) dose series may be used to meet this requirement. If you are unsure if a particular child's two (2) dose schedule is acceptable, please contact the Immunization Section for assistance at 501-661-2169.

*** 3rd dose of hepatitis B should be given at least eight (8) weeks after the 2nd dose, at least sixteen (16) weeks after the 1st dose, and it should not be administered before the child is twenty-four (24) weeks (one hundred and sixty-eight (168) days) of age. (All 3rd doses of hepatitis B vaccine given earlier than six (6) months of age before 06/21/1996 are valid doses and should be counted as valid until 06/21/2014.)

**** Tdap vaccine can be administered regardless of the interval since the last tetanus and diphtheria toxoid-containing vaccine.

***** Exception: If a student has previously received two (2) doses of measles, one (1) dose of mumps and one (1) dose of rubella before January 1, 2010, the doses will be accepted as compliant to immunization requirements and two (2) MMRs are not required.

*****A medical professional is a Medical Doctor (MD), Advanced Practice Nurse (APN), Doctor of Osteopathy (DO), or Physician Assistant (PA). No self or parental history of disease will be accepted.

***** For unvaccinated persons seven (7) years of age and older (including persons who cannot document prior vaccination), the primary series is three (3) doses. The first two (2) doses should be separated by at least four (4)

weeks, and the third dose at least six (6) months after the second. One (1) of these doses (preferably the first) should be administered as Tdap and the remaining two (2) doses administered as Td.